A set is a collection of definite well defined objects.

A set is a collection of objects which are distinct from each other.

A set is usually denoted by capital letter, i.e, A, B, S, T, G etc. A set elements are denoted by small letter, i.e, a, b, s, t etc.

CONSTRUCTION OF SET:

In construction of set, two methods are commonly used-

 Roster Method (Enumeration): In this method we prepare a list of objects forming the set, writing the elements one after another between a pair of curly brackets.
 For example:

 $A = \{a, b, c, d\}.$

2) Description Method: In this method we describe the set in symbolic language. For example:

A set of integer numbers which is divisible by 3 is written as,

 $A = \{x : x \text{ is an integer divisible by 3}\}$

TYPES OF SET:

1) Singleton set: If a set consisting only 1 element is known as singleton set.

For example:

 $A = \{a\}.$

2) Finite set: If a set consisting finite number of elements is known as finite set. For example: $A = \{2, 4, 6, 8\}.$

3) Infinite set: If a set consisting infinite number of elements is known as infinite set. For example-

The set of all natural numbers.

 $A = \{1, 2, 3, \dots\}$

4) Equal sets: Two sets A and B consisting of the same elements is known as equal set. For example:

 $A = \{a, b, c, d\}$ and $B = \{a, b, c, d\}$

5) Empty set: If a set consisting no elements is known as empty set or null set or void set. For example:

 $\mathsf{A} = \{ \emptyset \}$

6) Subset: Suppose A is a given set, and any set B exist exist whose elements are also an element of A,than B is called subset of A.
For example:
A = {1, 2, 3, 4, 5, 6, 7, 8} and B = {2, 4, 6, 8}
Than, B ⊆ A. (read as B is the subset of A)

Now take another example;

A = {1, 2, 3, 4, 5, 6, 7, 8} and B = {1, 2, 3, 4, 5, 6, 7, 8} Than, B \subseteq A. (read as B is the subset of A)

7) Proper Subset: If B is the subset of A, and $B \neq A$, then B is proper subset of A. For example: A = {1, 2, 3, 4, 5, 6, 7, 8} and B = {2, 4, 6, 8} Than, B \subset A. (read as B is the proper subset of A)

8) Power set: The set of all subset of a set A, is known as power set of A.
For example:
A = {a, b, c}
Than

Power set, $P(A) = \{ \{ \emptyset \}, \{a\}, \{b\}, \{c\}, \{d\}, \{ab\}, \{ac\}, \{ad\}, \{bc\}, \{bd\}, \{cd\}, \{abc\} \}$

Related posts:

- 1. Mathematical induction
- 2. Relation
- 3. Net 34
- 4. prove that- $AX(B \cap C) = (AXB) \cap (AXC)$
- 5. Prove that- An(BuC) = (AnB) u (AnC)
- 6. prove that $-(A\cap B)X(C\cap D) = (AXC)\cap(BXD)$
- 7. Show that- $(P \cap Q)X(R \cap S) = (PXR) \cap (QXS)$
- 8. Binary operations
- 9. Algebraic structure
- 10. Group
- 11. Show that (..., -4, -3, -2, -1, 0, 1, 2, 3, 4,...} is group
- 12. Show that a*b=b*a
- 13. if $a^*c = c^*a$ and $b^*c = c^*b$, then $(a^*b)^*c = c^*(a^*b)$
- 14. Undirected Graph and Incident Matrix
- 15. Prove the following by using the principle of mathematical induction for all $n \in N$, $1^3 + 2^3 + 3^3 + ... + n^3 = [n (n + 1)/2]^2$
- 16. Prove that $G = \{-1, 1, i, -i\}$ is a group under multiplication.

17. Hasse diagram for the "less than or equal to" relation on the set $S = \{ 0,1,2,3,4,5 \}$